

This patient group direction (PGD) must only be used by registered health professionals who have been named and authorised by their organisation to practice under it. The most recent and in date final signed version of the PGD should be used.

Patient Group Direction (PGD)

For the administration or supply of

Codeine Phosphate tablets 30mg

By registered health care professionals for

Acute, moderate to severe pain

Throughout the Manx Care and those contracted by the Manx Care where appropriate within practice

PGD NUMBER 108

1. Change history

| Version number | Change details | Date |
|-------------------|--|------------------------|
| 1 | Original PGD ratified | June 2021 |
| 2 | Minor word changes to ensure clarify of PGD and links added to Cautions and Adverse Effects, and reference to Manx Care Policy for Capacity added. Plus: Cautions updated Amendments make to Exclusion criteria CD schedule 5 added to Legal Category | Nov 2023 & May 2024 |
| | | |

2. Medicines practice guideline 2: Patient group directions

Refer to the relevant sections of NICE medicines practice guideline 2: *Patient group directions* as stated in the blank template notes. For further information about PGD signatories, see the NHS and Manx Care <u>PGD website FAQs</u>.

3. PGD development

Refer to the NICE PGD competency framework for people developing PGDs

| Job Title & organisation | Name | Signature | Date |
|---------------------------------|------|-----------|------|
| Author of the PGD | | | |
| Member of the PGD working group | | | |

4. PGD authorisation

Refer to the <u>NICE PGD competency framework for people authorising PGDs</u>

| Pre Signatures | | | |
|--|------|-----------|------|
| Job Title | Name | Signature | Date |
| Chief Pharmacist | | | |
| Head of Ambulance Services | | | |
| GP Adviser | | | |
| Senior Microbiologist (if PGD contains antimicrobials) | N/A | N/A | N/A |
| Final signatures | | | |
| Medical Director | | | |
| Director of Nursing | | | |

5. Training and competency of registered healthcare professionals, employed or contracted by the Manx Care, GP practice or Hospice

| | Requirements of registered Healthcare professionals working | |
|---------------------------|--|--|
| | under the PGD | |
| Qualifications and | Registered healthcare professionals, working within or | |
| professional registration | contracted by the Manx Care, GP practice or Hospice who are | |
| | permitted staff groups outlined within the current PGD policy | |
| | Pharmacists must be practising in Manx Care authorised | |
| | premises i.e. contracted pharmacy premises | |
| Initial training | Knowledge of current guidelines and the administration of the drug specified in this PGD/BNF and of the inclusion and exclusion criteria | |
| | • Training which enables the practitioner to make a clinical | |
| | assessment to establish the need for the medication covered by | |
| | this PGD | |
| | Local training in the use of PGDs | |
| Competency | Staff will be assessed on their knowledge of drugs and clinical | |
| assessment | assessment as part the competency framework for registered health | |
| | professionals using PGDs | |
| Ongoing training and | The registered health care professionals should make sure they are | |
| competency | aware of any changes to the recommendations for this medication; | |
| _ | it is the responsibility of the registered health care professionals to | |
| | keep up to date with continuing professional development. PGD | |
| | updates will be held every two years | |

Refer to the NICE PGD competency framework for health professionals using PGDs

6. Clinical Conditions

| Clinical condition or situation to which this PGD applies | Acute, moderate to severe pain |
|---|---|
| Inclusion criteria | Patients 12 years and overAcute moderate to severe pain |
| Exclusion criteria | Children under age 12 Acute respiratory depression Acute asthma/COPD Acute colitis Allergy and hypersensitivity to Codeine Antibiotic associated colitis Breastfeeding Conditions where abdominal distension develops Children under age 12 Conditions where peristalsis may be impaired Head injury/raised ICP Hypothyroidism Hypotension (systolic <100mmHg) Inflammatory or obstructive bowel disorders Risk of paralytic ileus Third trimester of pregnancy When using other opiate based medicines |

| Cautions (including any | Patients who are pregnant | |
|--|--|--|
| relevant action to be | Asthma/COPD | |
| taken) | Biliary tract disease | |
| | Cardiac arrhythmias | |
| | Chronic malnutrition or dehydration | |
| | | |
| | | |
| | Hepatocellular insufficiency History of alcohol dependence, or shronis alcoholism | |
| | History of alcohol dependence, or chronic alcoholism | |
| | History of drug addiction or those currently prescribed Naltrexone | |
| | Inflammatory bowel disorders, and obstructive bowel disorders | |
| | Marked increase in morphine toxicity in patients who are ultra- | |
| | rapid codeine metabolisers (CYP2D6 Ultra Rapid Metabolisers) | |
| | Myasthenia gravis | |
| | Prostatic hypertrophy | |
| | Reduced therapeutic effect in poor codeine metabolisers | |
| | Shock | |
| | Urethral stenosis | |
| | MHRA advice : Caution should be taken as to prescribing opioid analgesics in patients prescribed benzodiazepines | |
| | analgesics in patients prescribed benzodiazepines | |
| | | |
| | A detailed list of cautions is available in the SPC, which is available | |
| | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: | |
| Arrangements for referral | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: <u>www.medicines.org.uk</u> and BNF <u>https://bnf.nice.org.uk</u> | |
| Arrangements for referral for medical advice | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: | |
| - | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: <u>www.medicines.org.uk</u> and BNF <u>https://bnf.nice.org.uk</u> Patient should be referred to a more experienced clinical | |
| for medical advice | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: <u>www.medicines.org.uk</u> and BNF <u>https://bnf.nice.org.uk</u> Patient should be referred to a more experienced clinical practitioner for further assessment | |
| for medical advice Action to be taken if | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: <u>www.medicines.org.uk</u> and BNF <u>https://bnf.nice.org.uk</u> Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical | |
| for medical advice Action to be taken if patient excluded | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: <u>www.medicines.org.uk</u> and BNF <u>https://bnf.nice.org.uk</u> Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical practitioner for further assessment | |
| for medical advice Action to be taken if patient excluded Action to be taken if | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk and BNF <u>https://bnf.nice.org.uk</u> Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical practitioner for further assessment • A verbal explanation should be given to the patient on: the need | |
| for medical advice Action to be taken if patient excluded Action to be taken if patient declines | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk and BNF <u>https://bnf.nice.org.uk</u> Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical practitioner for further assessment A verbal explanation should be given to the patient on: the need for the medication and any possible effects or potential risks | |
| for medical advice Action to be taken if patient excluded Action to be taken if patient declines | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk and BNF https://bnf.nice.org.uk Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical practitioner for further assessment A verbal explanation should be given to the patient on: the need for the medication and any possible effects or potential risks which may occur as a result of refusing treatment | |
| for medical advice Action to be taken if patient excluded Action to be taken if patient declines | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk and BNF https://bnf.nice.org.uk Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical practitioner for further assessment A verbal explanation should be given to the patient on: the need for the medication and any possible effects or potential risks which may occur as a result of refusing treatment This information must be documented in the patients' health records Any patient who declines care must have demonstrated capacity | |
| for medical advice Action to be taken if patient excluded Action to be taken if patient declines | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk and BNF https://bnf.nice.org.uk Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical practitioner for further assessment A verbal explanation should be given to the patient on: the need for the medication and any possible effects or potential risks which may occur as a result of refusing treatment This information must be documented in the patients' health records Any patient who declines care must have demonstrated capacity to do so (see the Manx Care Policy for Capacity, Best Interests | |
| for medical advice Action to be taken if patient excluded Action to be taken if patient declines | A detailed list of cautions is available in the SPC, which is available from the electronic Medicines Compendium website: www.medicines.org.uk and BNF https://bnf.nice.org.uk Patient should be referred to a more experienced clinical practitioner for further assessment Patient should be referred to a more experienced clinical practitioner for further assessment A verbal explanation should be given to the patient on: the need for the medication and any possible effects or potential risks which may occur as a result of refusing treatment This information must be documented in the patients' health records Any patient who declines care must have demonstrated capacity | |

7. Details of the medicine

| Name, form and strength | Codeine Phosphate 30mg tablets |
|----------------------------|----------------------------------|
| of medicine | |
| Legal category | Prescription Only Medicine (POM) |
| | CD schedule 5 |
| Indicate any off-label use | None |
| (if relevant) | |

| Route/method of | Oral | |
|---------------------|---|--|
| administration | | |
| | 12 years and over 20 60mg every 6 hours if required (4.6 hours) | |
| Dose and frequency | 12 years and over: 30 – 60mg every 6 hours if required (4-6 hourly) | |
| Quantity to be | Administered: One by stated dose | |
| administered and/or | Supplied: One original pack | |
| supplied | | |
| Maximum or minimum | 12 years and over: maximum 240mg per day for maximum of | |
| treatment period | 3 days | |
| Storage | Room temperature | |
| Adverse effects | Arrhythmias Hypotension (with high doses) | |
| | Confusion Miosis | |
| | Constipation Nausea (more common on initiation) | |
| | Dizziness Palpitations | |
| | Drowsiness Respiratory depression (with high doses) | |
| | Dry mouth Skin reactions | |
| | Eeuphoric Urinary retention | |
| | mood • Vertigo | |
| | Flushing Visual impairment | |
| | Hallucination Vomiting (more common on initiation) | |
| | Headache Withdrawal syndrome | |
| | Hyperhidrosis | |
| | • Hypermatosis | |
| | A detailed list of adverse reactions is available in the SPC, which is available from the electronic Medicines Compendium website: <u>www.medicines.org.uk</u> and BNF <u>https://bnf.nice.org.uk</u> | |
| Records to be kept | The administration of any medication given under a PGD must be recorded within the patient's medical records | |
| | | |

8. Patient information

| Verbal/Written information to be given to patient or carer | Verbal information must be given to patients and or carers for all medication being administered under a PGD Where medication is being supplied under a PGD, written patient information leaflet must also be supplied A patient information leaflet is available on request Patients should be advised to avoid driving and or operating heavy machinery should this medication cause drowsiness | |
|--|--|--|
| Follow-up advice to be | If symptoms do not improve or worsen or you become unwell, seek | |
| given to patient or carer medical advice immediately | | |

9. Appendix A

References

- 1. British National Formulary (BNF) available online: <u>https://bnf.nice.org.uk</u>
- 2. Nursing and Midwifery "The code" available online: <u>https://www.nmc.org.uk</u>
- 3. Current Health Care Professions Council standards of practice
- 4. General Pharmaceutical Council standards
- 5. Electronic medicines compendium available online: <u>https://www.medicines.org.uk</u>
- 6. Manx Care Policy for Capacity, Best Interests Decisions and Deprivation of Liberty <u>http://edrmgi/sites/hospital/Clinical%20Policies%20and%20Procedures/Policy%20for%20Cap</u> <u>acity,%20Best%20Interests%20Decisions%20and%20Deprivation%20of%20Liberty.pdf#search</u> <u>=deprivation</u>

10. Appendix B

Health professionals agreed to practice

- Each registered healthcare professional will hold their own Competency framework which will be signed and agreed by their mentor
- A mentor is defined within the Manx Care policy as any ward/area managers, sisters, senior nurses, GPs, pharmacists or senior paramedics who has completed the PGD training themselves